

Link to the final version: <https://doi.org/10.1111/sms.13566>

**TITLE PAGE**

**Title**

Associations of 24-h activity composition with adiposity and cardiorespiratory fitness: the PregnActive Project.

**Running head**

24-h activity composition during pregnancy.

**Authors**

OVIEDO-CARO, Miguel Angel. [maovicar@upo.es](mailto:maovicar@upo.es) PhD <sup>1</sup>Physical Performance & Sports Research Center, Department of Sports and Computer Science, Section of Physical Education and Sports, Faculty of Sports Sciences, University Pablo de Olavide, Spain. <sup>2</sup>Research group in development Movimiento Humano. Universidad de Zaragoza, Spain.

BUENO-ANTEQUERA, Javier. [jbuant@upo.es](mailto:jbuant@upo.es) PhD <sup>1</sup>Physical Performance & Sports Research Center, Department of Sports and Computer Science, Section of Physical Education and Sports, Faculty of Sports Sciences, University Pablo de Olavide, Spain. <sup>2</sup>Research group in development Movimiento Humano. Universidad de Zaragoza, Spain.

MUNGUÍA-IZQUIERDO, Diego. [dmunizq@upo.es](mailto:dmunizq@upo.es) PhD <sup>1</sup>Physical Performance & Sports Research Center, Department of Sports and Computer Science, Section of Physical Education and Sports, Faculty of Sports Sciences, Universidad Pablo de Olavide, Seville, Spain. <sup>2</sup>Biomedical Research Networking Center on Frailty and Healthy Aging, Madrid, Spain. <sup>3</sup>Research group in development Movimiento Humano. Universidad de Zaragoza, Spain.

**Corresponding author:**

Oviedo-Caro, Miguel Ángel.

26 E-mail address: [maovicar@upo.es](mailto:maovicar@upo.es)

27 Twitter username: @Oviedo-CaroMA

28 Mailing address: University Pablo de Olavide, Carretera de Utrera Km 1, ES-41013

29 Seville, Spain

30 Telephone number: +34 954977589

31 Fax number: +34 954348377

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

**ABSTRACT**

**Aim**

This study examined the associations between activity behaviors composition (sleep, sedentary time, light and moderate-to-vigorous physical activity) with adiposity and cardiorespiratory fitness, and how isothermal reallocations of time between activity behaviors are associated with differences in adiposity and cardiorespiratory fitness.

**Methods**

A cross-sectional study was conducted in 130 women during midpregnancy. Activity behaviors, conceptualized as a 24-hour composition, were objectively assessed by multi-sensor monitors. Skinfold thickness, fat mass index, and body mass index were calculated as indicators of adiposity. Cardiorespiratory fitness was assessed using a 6-minute walk test. Log-ratio multiple linear regression models and compositional isothermal substitutions were used to analyze the associations and estimated differences in outcomes.

**Results**

The activity composition was significantly associated with adiposity indicators (all  $p < 0.001$ ) and cardiorespiratory fitness ( $p$  values from 0.025 to  $< 0.001$ ) during midpregnancy. The isothermal substitutions were asymmetrical, showing the highest estimated differences in adiposity (8.7%, 0.80 kg/m<sup>2</sup>, for fat mass index; 6.0%, 2.65 mm, for the sum of skinfold thickness; and 3.8%, 1.02 kg/m<sup>2</sup>, for body mass index) and cardiorespiratory fitness (3.0%, 1.00 mL/kg·min) when 30-minutes of moderate-to-vigorous physical activity was reallocated by sedentary time.

**Conclusion**

The activity composition was associated with adiposity and the cardiorespiratory fitness levels during midpregnancy, with moderate-to-vigorous physical activity being the

leading activity behavior. The most unfavorable differences in adiposity and cardiorespiratory fitness were found when moderate-to-vigorous physical activity was replaced by another behavior, mainly sedentary time, reinforcing the importance of at least maintaining moderate-to-vigorous physical activity during pregnancy.

## KEYWORDS

Time-use; physical activity; sedentary time; sleep; physical fitness; adiposity.

PROOF

1

2

31011. INTRODUCTION

4

5102Lifestyle physical activities have been proposed to be behaviors that may protect against

6

7103the development of cardiovascular diseases or prevent recurrent cardiovascular events <sup>1</sup>.

8

9

10104Especially during pregnancy, lifestyle physical activities have potential benefits to

11

12105reduce the risk of adverse cardiovascular health outcomes, such as hypertension, and

13

14106obesity <sup>2,3</sup>. Changes in the maternal body composition occur over a short period of time

15

16

17107during pregnancy, usually after the first trimester <sup>4</sup>. The increase of the mother's

18

19108adiposity is a serious problem because of its association with an elevated risk of

20

21109pregnancy-related complications, long-term maternal obesity and childhood obesity <sup>5,6</sup>.

22

23

24110Cardiorespiratory fitness (CRF) exhibits protective effects on established pregnancies

25

26111and pregnant women's cardio-metabolic health outcomes, such as insulin sensitivity and

27

28112glucose and lipid disposal <sup>7</sup>. However, limitations in aerobic work capacity, caused by

29

30113an increase in the ventilatory response related to the elevated metabolic costs of

31

32114exercise, result in a decline of CRF levels as pregnancy advances <sup>8</sup>.

33

34

35115Previous research has focused on the effect of the time spent in ST or MVPA

36

37116individually, on adiposity and on CRF, promoting MVPA and discouraging ST among

38

39117pregnant women <sup>9-12</sup>. Taking into account that activity behaviors are mutually exclusive

40

41118components of the 24-h day, this time has to be reallocated from another activity

42

43119behavior, which may have different effect on health outcomes depending on what

44

45120activity behavior is reallocated.

46

47

48

49121This 24-h activity behaviors concept is related to the notion that to properly understand

50

51122the association between daily activities and health, the effect of all activity behaviors

52

53123should be studied relative to each other rather than in isolation <sup>13,14</sup>. The application of

54

55124this concept implies that activity behaviors data are treated as compositional data that

56

57125are composed of mutually exclusive and exhaustive parts of a whole 24-h <sup>13</sup>.

58

59

60

To date, only two studies have applied compositional data analysis to assess the association of activity behaviors and adiposity in adults <sup>13,15</sup>, and only studies in children have assessed the association of activity behaviors with CRF <sup>14,16</sup>. No studies have applied compositional data analysis to pregnant women.

The aims of this study are (1) to assess the associations between the activity composition and adiposity and CRF during midpregnancy and (2) to investigate how time reallocations between activity behaviors are associated with favorable or unfavorable adiposity and CRF.

## **2. METHODS**

### **2.1 Participants and procedures**

This exploratory cross-sectional study was conducted in 130 healthy pregnant women aged 18-45 years who were recruited from antenatal clinics at Utrera Hospital, and voluntarily gave their written informed consent after being informed of the study aims and protocol. The exclusion criteria were physical illnesses or disabilities that affected their normal daily routine or high-risk pregnancy (i.e., diabetes or hypertension). The study protocol obtained ethical approval from the Medical Research Ethics Committee of the University Hospital Virgen del Rocío (Seville, Spain) in accordance with the Declaration of Helsinki, approval number 2014PI-066. The STROBE guideline items were fulfilled during the course of the study <sup>17</sup>.

Coinciding with the second visit to the antenatal clinics, approximately at the 20<sup>th</sup> gestational week, the sociodemographic characteristics, adiposity, and CRF were assessed.

### **2.2 Measurements**

#### **2.2.1 Sociodemographic characteristics**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

150 Age, educational level and employment status were assessed using a self-report  
151 questionnaire. The pregnant women were categorized according to having a tertiary or  
152 non-tertiary educational level, and active or non-active (unemployed, sick-leave from  
153 work and student) employment status.

154 **2.2.2 Adiposity**

155 Anthropometry was measured by the same trained evaluator using a stadiometer to  
156 measure height, a bioelectrical impedance analysis (BIA) device (Tanita BC-420,  
157 Tanita, Tokyo, Japan) to measure weight and the body composition, and a caliper  
158 (Holtain, Crymich, UK) to measure skinfold thickness. The sum of the biceps, triceps,  
159 and subscapular skinfold thicknesses, measured to the nearest millimeter on the right-  
160 hand side of the body, was used as an indicator of maternal adiposity as previously  
161 proposed <sup>5,18</sup>. For comparison with previous studies, body mass index (BMI, kg/m<sup>2</sup>) and  
162 fat mass index (FMI, kg/m<sup>2</sup>) were calculated.

163 **2.2.3 Cardiorespiratory fitness**

164 CRF was assessed using a 6-minute walk test (6MWT), previously used with pregnant  
165 women <sup>9,19</sup>, through a 45.7-meter rectangular course delimited by cones. Participants  
166 were encouraged to walk as far as possible without running or jogging. The same  
167 trained instructor explained the protocol, provided a demonstration prior to the start,  
168 supervised the test and recorded the distance covered to the nearest 0.1 m. A multimedia  
169 explanation is available on the link below:  
170 <https://upotv.upo.es/video/5936500f238583f9658b464a>. The resting heart rate was  
171 monitored beat-to-beat after a 5-minute seated rest period using a heart rate monitor  
172 (Polar Electro Oy, Kempele, Finland). Maximal oxygen consumption (VO<sub>2max</sub>,  
173 mL/kg·min) was estimated from the 6MWT using the equation reported for healthy  
174 adults by Burr et al. <sup>20</sup>.

#### 2.2.4 24-h activity behaviors

Free-living activity behaviors were objectively measured using a multi-sensor monitor Sensewear Mini Armband (BodyMedia Inc., Pittsburgh, PA, USA) (SWA), validated in pregnant women <sup>21,22</sup>, over a 24-hours during a 9-day period, including at least five weekdays and two weekend days. Participants were told to remove the monitor only for water-based activities, which were recorded in a diary. Only participants who carried the monitor for at least 95% of the day (1368 minutes) were included in the study. Five participants participated in swimming and water aerobics or calisthenics for a maximum of 45 minutes, and the time spent in these activities was added using a constant (6 and 4 METs, respectively) from the Compendium of Physical Activities <sup>23</sup>. In addition, non-wear time from showering and self-care were substituted using a constant (2 METs) from the Compendium. The sleep component represents all sleep occurring between 12 PM and 12 AM. The sleep component did not necessarily describe the overnight sleep duration, and it did incorporate naps taken during day. To minimize immediate reactivities that may have altered their habitual lifestyle, we removed the first and the last day of monitoring from the analysis.

#### 2.3 Statistical analysis

Analysis were performed in R using the *compositions* <sup>24</sup>, *robcompositions* <sup>25</sup> and *car* <sup>26</sup> packages. The average daily time spent in free-living activity behaviors (sleep, ST, LPA and MVPA) was conceptualized as part of the daily activity behaviors composition and was expressed as isometric log-ratio (ilr) coordinates <sup>13</sup>. All free-living activity behaviors contain non-zero values. The multi-variate dispersion of the activities was described by a pair-wise variation matrix.

Linear regression models were fitted to investigate the associations between adiposity and CRF as outcome variables, and the activity behaviors' ilr coordinate as explanatory



1  
2  
3 200 variables. Sociodemographic covariates were also included as explanatory variables.  
4  
5 201 The significance of the explanatory variables was examined by a Wald chi square  
6  
7 202 ANOVA type II test <sup>26</sup>. In all models, non-normally distributed variables (sum of  
8  
9 203 skinfold thickness, FMI, and BMI) were log-transformed. Residuals were tested for  
10  
11 204 normality, linearity, homoscedasticity, and independence.  
12  
13  
14 205 Compositional isotemporal substitutions were used to examine the difference in the  
15  
16 206 outcome variables when quanta of time (from 5 to 30 minutes) were reallocated from  
17  
18 207 one activity to another, while the remaining activities were kept constant <sup>27</sup>. The above  
19  
20 208 models were used as predictive models for new activity behaviors compositions. First,  
21  
22 209 the model was used to predict an outcome value for a baseline composition (mean  
23  
24 210 composition), and subsequently, the same model was used to predict the outcome for  
25  
26 211 new compositions when quanta of time had been reallocated. The estimated differences  
27  
28 212 in the outcome variable were calculated by subtracting the predicted values of the mean  
29  
30 213 composition from the predicted values of the new composition. The significance of the  
31  
32 214 reallocation was assessed based on 95% CI as previously suggested <sup>16</sup>. Effect sizes (ES)  
33  
34 215 were calculated by dividing the estimated differences by the mean standard deviation of  
35  
36 216 outcome variables as previously suggested <sup>28</sup>.

41  
42 217 **3. RESULTS**

43  
44 218 The study sample was composed of 130 pregnant women, after excluding 36  
45  
46 219 participants with incomplete data, and their characteristics and compositional means for  
47  
48 220 their activity behaviors are presented in the Supplemental Table 1. The compositional  
49  
50 221 variation matrix is presented in Supplemental Table 2. The smallest variances were  
51  
52 222 observed for sleep with ST and LPA, and ST with LPA (values  $\leq 0.195$ ), indicating the  
53  
54  
55 223 greatest co-dependence between activities.  
56  
57  
58  
59  
60

Multiple linear regression models showed that the activity composition was significantly associated with all adiposity indicators and CRF (Table 1). The predicted outcome values for the mean composition were 44.3 mm for the sum of skinfold thickness, 9.1 kg/m<sup>2</sup> for FMI, 26.9 kg/m<sup>2</sup> for BMI, and 33.8 mL/kg·min for CRF.

Compositional isotemporal substitutions were carried out for adiposity indicators and CRF. The raw estimated differences and 95% CI in adiposity and CRF associated with increments of time reallocations are shown in Supplemental Tables 3 and 4, respectively. For a better interpretation of the results, Tables 2 and 3 show the estimated percentage of difference from the mean composition in predicted adiposity and CRF associated with 15-minute and 30-minute reallocations. The reallocations of time involving MVPA was associated with the largest estimated differences in adiposity and CRF and are plotted in Supplemental Figure 1.

The largest estimated differences found in adiposity were observed for the reallocation of time from MVPA to ST. Among the differences the largest differences were observed for FMI, followed by the sum of skinfold thickness, and BMI, although in a small magnitude. For instance, reallocation of 30-minutes from MVPA to ST was associated with higher adiposity, approximately 8.7% for FMI (0.80 kg/m<sup>2</sup>, 95% CI: 0.33 to 1.29, ES: 0.23), 6.0% for the sum of skinfold thickness (2.65 mm, 95% CI: 0.83 to 4.52, ES: 0.21), and 3.8% for BMI (1.02 kg/m<sup>2</sup>, 95% CI: 0.42 to 1.64, ES: 0.25) compared with the mean composition. The association showed marked asymmetry, the estimated difference in adiposity associated with the reallocation of time from ST to MVPA was lower than the aforementioned estimated difference in adiposity associated with the reallocation of time from MVPA to ST. For instance, reallocation of 30-minutes from ST to MVPA was associated with a difference of -6.1% for FMI (-0.56 kg/m<sup>2</sup>, 95% CI: -0.84 to -0.27, ES: -0.17), -4.2% for the sum of skinfold thickness (-1.88 mm, 95% CI: -

1  
2  
3 249 3.00 to -0.73, ES: -0.13), and -2.8% for BMI (-0.76 kg/m<sup>2</sup>, 95% CI: -1.14 to -0.36, ES: -  
4  
5 250 0.17).  
6  
7  
8 251 The largest estimated differences in CRF were observed for the reallocation of time  
9  
10 252 from MVPA to ST or sleep. For instance, the reallocation of 30-minutes was associated  
11  
12 253 with a lower CRF by approximately 4.0% (-1.39 mL/kg·min, 95% CI: -2.00 to -0.78,  
13  
14 254 ES: -0.29, and -1.35 mL/kg·min, 95% CI: -2.08 to -0.62, ES: -0.28, respectively)  
15  
16  
17 255 compared with the mean composition. The association showed a marked asymmetry,  
18  
19 256 the estimated difference in CRF associated with a 30-minute reallocation from ST and  
20  
21 257 sleep to MVPA, 3.0% of the mean composition (1.00 mL/kg·min, 95% CI: 0.58 to 1.41,  
22  
23 258 ES: 0.21, and 0.97 mL/kg·min, 95% CI: 0.42 to 1.52, ES: 0.20, respectively), was lower  
24  
25  
26 259 than the aforementioned estimated difference in CRF associated with the inverse  
27  
28 260 reallocation of time. When the model was additionally adjusted for adiposity, the  
29  
30 261 magnitudes of the estimated differences were lower, especially when the model was  
31  
32 262 adjusted for FMI, or the reallocations implying ST and MVPA when the model was  
33  
34 263 adjusted for BMI.  
35  
36

37 264 **4. DISCUSSION**

38  
39  
40 265 Application of the novel compositional data analysis approach presented in this study  
41  
42 266 allows for a proper analysis of the influence of activity behaviors as mutually exclusive  
43  
44 267 components of the 24-h day on adiposity and CRF. Pregnant women's activity  
45  
46 268 composition was significantly associated with all of the adiposity indicators and CRF,  
47  
48 269 with MVPA being the activity behavior with the greatest association. Isotemporal  
49  
50 270 substitution models suggested that the most unfavorable differences in adiposity and  
51  
52 271 CRF were when MVPA was replaced by ST. Estimated differences were relatively  
53  
54  
55 272 modest, with standardized effect sizes ranging between 0.2 and 0.3. This result  
56  
57  
58  
59  
60

reinforces the importance of at least maintaining MVPA during pregnancy, particularly when taking into account the usual decrease of MVPA levels as pregnancy advances <sup>29</sup>. Consistent with previous compositional isothermal substitution studies in adults <sup>13,15</sup>, the highest predicted differences in adiposity indicators were found when MVPA was reallocated to ST. The estimated differences of the sum of skinfold thickness differed for FMI and BMI, which could be explained by their inherent limitations in differentiating between maternal and fetal contributions for adiposity. BMI showed the lowest estimated differences, which could be explained by BMI accounting for lean mass as well, which may be higher while adiposity may be lower with MVPA <sup>30</sup>. Previous studies on pregnant women have shown how MVPA, such as walking and exercise interventions, is associated with the control of maternal adiposity, lipid biomarkers, and weight gain <sup>10,11</sup>, but as our results suggest, the greatest benefits of differences in adiposity were obtained when ST was reallocated to MVPA. The fact that physical activity was associated with a lower adiposity could be explained by the ability of exercise to moderate glucose levels by increasing glucose uptake of skeletal muscle and improving insulin sensitivity <sup>31</sup>. Future interventions aimed at maintaining MVPA levels or reducing ST in favor of MVPA may be effective ways to preserve adiposity levels among pregnant women.

The improvement of CRF by increasing MVPA levels is consistent with a previous compositional data study performed on children <sup>14,16</sup> and an isothermal non-compositional substitution study performed on the general population <sup>32</sup>. In addition, randomized controlled trials in pregnant women <sup>9,11</sup> have suggested that an increase in MVPA levels may improve pregnant women's CRF, but as our results suggest, the highest benefit on CRF was obtained when ST, instead of others behaviors, was reallocated to MVPA. Although the reallocation of time from sleep to MVPA is

1  
2  
3 298 associated with a similar difference in CRF, considering the important role of sleep in  
4  
5 299 pregnant women's health and the fact that pregnancy alters sleep patterns <sup>33</sup>, it would be  
6  
7 300 inappropriate to propose it when pregnant women do not fulfill sleep recommendations  
8  
9  
10 301 <sup>34</sup>. Reallocating time from MVPA to ST was associated with the most unfavorable  
11  
12 302 differences in CRF. Taking into account that usually ST increases and MVPA decreases  
13  
14 303 along the course of pregnancy <sup>35</sup>, our results suggest that interventions aimed to at least  
15  
16 304 maintain MVPA and ST levels, or reduce ST in favor of MVPA, may be recommended  
17  
18 305 to limit the decrease in CRF caused by the progression of pregnancy <sup>8</sup>.  
19  
20  
21 306 The associations of the reallocation of time with adiposity and CRF, consistent with  
22  
23 307 previous compositional data studies <sup>13,16</sup>, are asymmetrical. For instance the reallocation  
24  
25 308 of time from MVPA to ST led to larger estimated differences in adiposity and CRF than  
26  
27 309 the reallocation of time from ST to MVPA. This difference could be explained by the  
28  
29 310 relative contribution of the different behaviors to the 24-h period. ST accounted for 42%  
30  
31 311 of the day compared with 22% for LPA or 5% for MVPA; therefore, taking 15 minutes  
32  
33 312 from MVPA or LPA is more relevant than taking 15 minutes from ST. In addition, as  
34  
35 313 previous study suggested <sup>13</sup>, increasing adiposity or deconditioning occurs rapidly when  
36  
37 314 activity levels drop, while returning to the previous level requires a larger amount of  
38  
39 315 exercise. Therefore, the results highlight the importance of maintaining the levels of  
40  
41 316 MVPA to preserve adiposity and CRF levels during pregnancy, a period of time along  
42  
43 317 which these outcomes unfavorably change <sup>4,8</sup>.  
44  
45  
46  
47  
48  
49 318 The results of this study reinforce the public message suggesting the development of  
50  
51 319 preventive health strategies that focus on the reduction of ST by increasing moving. The  
52  
53 320 fact that ST is the most prevalent behavior among pregnant women during  
54  
55 321 midpregnancy <sup>36</sup>, and increases from pre-pregnancy <sup>35</sup> and also from midpregnancy to  
56  
57 322 later pregnancy (about 30 minutes) <sup>37</sup>, supports the importance of strategies aimed at  
58  
59  
60

reducing ST among pregnant women. As our results suggest, ST may be reallocated to MVPA to obtain the greatest benefits. However, the increase of MVPA during pregnancy may be difficult due to the increases in basal energy expenditure and the elevated metabolic costs of exercise during pregnancy<sup>8</sup>, which explain women's selection of less-demanding activities<sup>38</sup> during pregnancy. Therefore, as our results showed, LPA, which accounts for 80% of the pregnant women's time spent in physical activity<sup>39</sup>, may be a good way or a progressive step to reallocate ST, although with lower benefits than those obtained by MVPA. This is an interesting finding because the reallocation involving LPA in others CODA studies did not show this association with health benefits<sup>13,16</sup>, and as we aforementioned could be explained with the particular characteristics of pregnant women, which could obtain health benefits replacing ST with LPA.

The contribution of physical activity to maintaining general fitness levels has been shown even when physical activity is only started during pregnancy<sup>40</sup>. Considering that pregnancy is identified as a teachable moment in women's life, in which the attention to both women's and fetus' health increases, this stage of life is thought to offer an ideal opportunity to target interventions introducing behavioral changes<sup>41</sup>.

#### 4.1 Strengths

This study is the first to apply the principles of a novel compositional data approach during pregnancy to analyze the influence of activity behaviors relative to each other on pregnant women's adiposity and CRF. The use of a multi-sensor monitor provides an objective and direct estimation of all activity behaviors based on energy expenditure, allowing a precise measure of 24-h activity data. Strict analysis of the data, including only participants who carried the monitor for at least 95% of the entire day (mean wear time of 1421 min/day) and removing the first and last day of monitoring, guaranteed the

1  
2  
3 348 representativeness of the daily activity measure. Adiposity was assessed according the  
4  
5 349 sum of the biceps, triceps and subscapular skinfold thicknesses, which has been  
6  
7  
8 350 proposed to reasonably estimate maternal fat mass and changes in maternal adipose  
9  
10 351 tissue that are not influenced by fetal growth <sup>5,18</sup>. For comparison with previous studies  
11  
12 352 we also use BIA's FMI and BMI, although we are conscious about their inherent  
13  
14 353 limitations during pregnancy.

16  
17 354 **4.2 Limitations**

18  
19 355 This study has several limitations. First, the cross-sectional design precludes the  
20  
21 356 establishment of any causal associations. Future studies should analyze the effects of  
22  
23 357 applying interventions that replace ST with MVPA or LPA, to confirm the causal  
24  
25 358 effects for individuals. The fact that high-risk pregnant women were excluded from the  
26  
27 359 study sample and the use of voluntary participation could have resulted in self-selection  
28  
29 360 bias. Future studies with larger sample sizes are required. Multi-sensor monitors cannot  
30  
31 361 differentiate between body positions, such as standing and ST. Studies that combine  
32  
33 362 multi-sensor monitors with postural monitors would better estimate ST. The use of FMI  
34  
35 363 or BMI has inherent limitations related to the impossibility of differentiating between  
36  
37 364 maternal and fetal contributions or between body fat and lean mass, respectively. A  
38  
39 365 submaximal test was used to estimate  $VO_{2max}$  although it has inherent limitations, the  
40  
41 366 6MWT has an acceptable coefficient of variation compared with direct measures <sup>20</sup>.  
42  
43 367 Future studies using a direct measure of  $VO_{2max}$  would improve our design. In addition,  
44  
45 368 the development and validation of a specific equation to estimate  $VO_{2max}$  for pregnant  
46  
47 369 women could be recommended for future studies on this population.

50  
51 370 **5. CONCLUSION**

52  
53 371 The activity composition was associated with adiposity and CRF levels during  
54  
55 372 midpregnancy, with MVPA being the leading activity behavior. Estimated differences  
56  
57  
58  
59  
60



were relatively modest, showing the most unfavorable differences in adiposity and CRF levels when MVPA was replaced mainly by ST. The findings reinforce the importance of at least maintaining MVPA and limiting the time spent in ST during pregnancy.

## 6. PERSPECTIVE

The findings of the present study might be considered relevant for clinical practice. Our results contribute to a better understanding of the associations between 24-h activity behaviors and health outcomes during pregnancy, providing some novelties to the scientific literature, with the application of the principles of the novel compositional data approach during pregnancy. Previous study suggested the usefulness of promoting MVPA and discouraging ST during pregnancy <sup>42</sup>, but studying activity behaviors in isolation may be misleading, even more so when the fulfillment of MVPA recommendations is not necessarily related with a reduction of sedentary time during midpregnancy <sup>29</sup>. Consequently, the reallocation of time from ST to MVPA or LPA, although with lower benefits, could be an efficient way to preserve or improve adiposity and CRF levels at midpregnancy. Moreover, future research is warranted to analyze the effects of interventions reallocating ST to MVPA or LPA during midpregnancy.

## ACKNOWLEDGEMENTS

The authors gratefully acknowledge all women for their collaboration. We also acknowledge the antenatal health clinic members from Utrera Hospital involved in the recruitment for their effort and great enthusiasm. The study was funded by Research Group CTS-948 of University Pablo de Olavide, Seville, Spain and European University of Madrid, Cátedra Real Madrid, Spain (funding project number P2017/RM08), this funding included material and equipment for the research. There was no external financial support. M.A.O. is supported by the Departamento de Innovación, Investigación y Universidad del Gobierno de Aragón y el Fondo Europeo



de Desarrollo Regional – Programa Operativo FEDER Aragón 2014-2020 “Construyendo Europa desde Aragón” (reference number PUI/2018-336). J.B. is supported by the Spanish Ministry of Education (grant number FPU13/05130) and by the Departamento de Innovación, Investigación y Universidad del Gobierno de Aragón y el Fondo Europeo de Desarrollo Regional – Programa Operativo FEDER Aragón 2014-2020 “Construyendo Europa desde Aragón” (reference number PUI/2018-337).

REFERENCES

1. Ozemek C, Laddu DR, Lavie CJ, Claeys H, Kaminsky LA, Ross R, Wisloff U, Arena R, Blair SN. An Update on the Role of Cardiorespiratory Fitness, Structured Exercise and Lifestyle Physical Activity in Preventing Cardiovascular Disease and Health Risk. *Prog Cardiovasc Dis* 2018;61:484–490.

2. da Silva SG, Ricardo LI, Evenson KR, Hallal PC. Leisure-Time Physical Activity in Pregnancy and Maternal-Child Health: A Systematic Review and Meta-Analysis of Randomized Controlled Trials and Cohort Studies. *Sports Med* 2017;47:295–317.

3. American College of Obstetricians and Gynecologists. Physical Activity and Exercise During Pregnancy and the Postpartum Period. *Obstet Gynecol* 2015;126:1326–1327.

4. Fattah C, Farah N, Barry SC, O’Connor N, Stuart B, Turner MJ. Maternal weight and body composition in the first trimester of pregnancy. *Acta Obstet Gynecol Scand* 2010;89:952–955.

5. Tomedi LE, Simhan HN, Chang CCH, McTigue KM, Bodnar LM. Gestational weight gain, early pregnancy maternal adiposity distribution, and maternal hyperglycemia. *Matern Child Health J* 2014;18:1265–1270.

6. Hull HR, Dinger MK, Knehans AW, Thompson DM, Fields DA. Impact of

- maternal body mass index on neonate birthweight and body composition. *Am J Obstet Gynecol* 2008;198:416.e1–6.
7. Thorell E, Goldsmith L, Weiss G, Kristiansson P. Physical fitness, serum relaxin and duration of gestation. *BMC Pregnancy Childbirth* 2015;15:168.
8. Davenport MH, Steinback CD, Mottola MF. Impact of pregnancy and obesity on cardiorespiratory responses during weight-bearing exercise. *Respir Physiol Neurobiol* 2009;167:341–347.
9. Ramirez-Velez R, Aguilar de Plata AC, Escudero MM, Echeverry I, Ortega JG, Salazar B, Rey JJ, Hormiga C, Lopez-Jaramillo P. Influence of regular aerobic exercise on endothelium-dependent vasodilation and cardiorespiratory fitness in pregnant women. *J Obs Gynaecol Res* 2011;37:1601–1608.
10. Ramírez-Vélez R, Lobelo F, Aguilar-de Plata AC, Izquierdo M, García-Hermoso A. Exercise during pregnancy on maternal lipids: A secondary analysis of randomized controlled trial. *BMC Pregnancy Childbirth* 2017;17:396.
11. Seneviratne SN, Jiang Y, Derraik JGB, McCowan LME, Parry GK, Biggs JB, Craigie S, Gusso S, Peres G, Rodrigues RO, Ekeroma A, Cutfield WS, Hofman PL. Effects of antenatal exercise in overweight and obese pregnant women on maternal and perinatal outcomes: A randomised controlled trial. *BJOG An Int J Obstet Gynaecol* 2016;123:588–597.
12. Mottola MF. Components of exercise prescription and pregnancy. *Clin Obstet Gynecol* 2016;59:552–558.
13. Chastin SFM, Palarea-Albaladejo J, Dontje ML, Skelton DA. Combined Effects of Time Spent in Physical Activity, Sedentary Behaviors and Sleep on Obesity and Cardio-Metabolic Health Markers: A Novel Compositional Data Analysis Approach. *PLoS One* 2015;10:e0139984.

1  
2  
3 448 14. Dumuid D, Stanford TE, Martin-Fernández JA, Pedišić Ž, Maher CA, Lewis LK,  
4  
5 449 Hron K, Katzmarzyk PT, Chaput JP, Fogelholm M, Hu G, Lambert E V, Maia J,  
6  
7 450 Sarmiento OL, Standage M, Barreira T V, Broyles ST, Tudor-Locke C, Tremblay  
8  
9 451 MS, Olds T. Compositional data analysis for physical activity, sedentary time  
10  
11 452 and sleep research. *Stat Methods Med Res* 2018;27:3726–3738.  
12  
13  
14 453 15. McGregor DE, Palarea-Albaladejo J, Dall PM, Stamatakis E, Chastin SFM.  
15  
16 454 Differences in physical activity time-use composition associated with  
17  
18 455 cardiometabolic risks. *Prev Med Reports* 2019;13:23–29.  
19  
20  
21 456 16. Fairclough SJ, Dumuid D, Mackintosh KA, Stone G, Dagger R, Stratton G,  
22  
23 457 Davies I, Boddy LM. Adiposity, fitness, health-related quality of life and the  
24  
25 458 reallocation of time between children’s school day activity behaviours: A  
26  
27 459 compositional data analysis. *Prev Med Reports* 2018;11:254–261.  
28  
29  
30 460 17. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP.  
31  
32 461 The Strengthening the Reporting of Observational Studies in Epidemiology  
33  
34 462 (STROBE) statement: guidelines for reporting observational studies. *J Clin*  
35  
36 463 *Epidemiol* 2008;61:344–349.  
37  
38  
39 464 18. Kannieappan LM, Deussen AR, Grivell RM, Yelland L, Dodd JM. Developing a  
40  
41 465 tool for obtaining maternal skinfold thickness measurements and assessing inter-  
42  
43 466 observer variability among pregnant women who are overweight and obese.  
44  
45 467 *BMC Pregnancy Childbirth* 2013;13:42.  
46  
47  
48 468 19. Oviedo-Caro MA, Bueno-Antequera J, Munguia-Izquierdo D. Explanatory  
49  
50 469 factors and levels of health-related quality of life among healthy pregnant women  
51  
52 470 at midpregnancy: A cross-sectional study of The PregnActive Project. *J Adv*  
53  
54 471 *Nurs* 2018;74:2766–2776.  
55  
56  
57 472 20. Burr JF, Bredin SSD, Faktor MD, Warburton DER. The 6-minute walk test as a  
58  
59  
60

- 473 predictor of objectively measured aerobic fitness in healthy working-aged adults.  
 474 Phys Sportsmed 2011;39:133–139.
- 475 21. Smith KM, Lanningham-Foster LM, Welk GJ, Campbell CG. Validity of the  
 476 SenseWear armband to predict energy expenditure in pregnant women. Med Sci  
 477 Sports Exerc 2012;44:2001–2008.
- 478 22. Conway MR, Marshall MR, Schlaff RA, Pfeiffer KA, Pivarnik JM. Physical  
 479 Activity Device Reliability and Validity during Pregnancy and Postpartum. Med  
 480 Sci Sports Exerc 2018;50:617–623.
- 481 23. Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DR, Tudor-Locke  
 482 C, Greer JL, Vezina J, Whitt-Glover MC, Leon AS. 2011 Compendium of  
 483 Physical Activities: a second update of codes and MET values. Med Sci Sport  
 484 Exerc 2011;43:1575–1581.
- 485 24. van den Boogaart KG, Tolosana-Delgado R. ‘compositions’: A unified R  
 486 package to analyze compositional data. Comput Geosci 2008;34:320–338.
- 487 25. Templ M, Hron K, Filzmoser P. 2011. robCompositions: An R-package for  
 488 Robust Statistical Analysis of Compositional Data. In: Compositional Data  
 489 Analysis: Theory and Applications.341–355.
- 490 26. Fox J, Weisberg S. An R Companion to Applied Regression. Thousand Oaks,  
 491 CA: SAGE; 2011.
- 492 27. Dumuid D, Pedišić Ž, Stanford TE, Martín-Fernández JA, Hron K, Maher CA,  
 493 Lewis LK, Olds T. The compositional isotemporal substitution model: A method  
 494 for estimating changes in a health outcome for reallocation of time between  
 495 sleep, physical activity and sedentary behaviour. Stat Methods Med Res 2017;
- 496 28. Dumuid D, Stanford TE, Pedišić Ž, Maher C, Lewis LK, Martín-Fernández JA,  
 497 Katzmarzyk PT, Chaput JP, Fogelholm M, Standage M, Tremblay MS, Olds T.

1  
2  
3 498 Adiposity and the isothermal substitution of physical activity, sedentary time  
4  
5 499 and sleep among school-aged children: A compositional data analysis approach.  
6  
7 500 BMC Public Health 2018;18:311.  
8  
9  
10 501 29. Di Fabio DR, Blomme CK, Smith KM, Welk GJ, Campbell CG. Adherence to  
11  
12 502 physical activity guidelines in mid-pregnancy does not reduce sedentary time: an  
13  
14 503 observational study. Int J Behav Nutr Phys Act 2015;12:27.  
15  
16  
17 504 30. Hueston WJ, Kasik-Miller S. Changes in functional health status during normal  
18  
19 505 pregnancy. J Fam Pract 1998;47:209–212.  
20  
21  
22 506 31. Hawley JA, Lessard SJ. Exercise training-induced improvements in insulin  
23  
24 507 action. Acta Physiol 2008;192:127–135.  
25  
26 508 32. Van Der Velde JHPM, Koster AM, Van der Berg JD, Sep SJS, Van der Kallen  
27  
28 509 CJH, DAGNELIE PC, Schram MT, Henry RMA, Eussen SJPM, Van Dongen  
29  
30 510 MCJM, Stehouwer CDA, Schaper NC, Savelberg HHCM. Sedentary Behavior,  
31  
32 511 Physical Activity, and Fitness—The Maastricht Study. Med Sci Sport Exerc  
33  
34 512 2017;49:1583–1591.  
35  
36  
37 513 33. Facco FL, Kramer J, Ho KH, Zee PC, Grobman WA. Sleep disturbances in  
38  
39 514 pregnancy. Obstet Gynecol 2010;115:77–83.  
40  
41  
42 515 34. Watson NF, Badr MS, Belenky G, Bliwise DL, Buxton OM, Buysse D, Dinges  
43  
44 516 DF, Gangwisch J, Grandner MA, Kushida C, Malhotra RK, Martin JL, Patel SR,  
45  
46 517 Quan SF, Tasali E, Twery M, Croft JB, Maher E, Barrett JA, Thomas SM, Heald  
47  
48 518 JL. Joint Consensus Statement of the American Academy of Sleep Medicine and  
49  
50 519 Sleep Research Society on the Recommended Amount of Sleep for a Healthy  
51  
52 520 Adult: Methodology and Discussion. J Clin Sleep Med 2015;11:931–952.  
53  
54  
55 521 35. Fazzi C, Saunders DH, Linton K, Norman JE, Reynolds RM. Sedentary  
56  
57 522 behaviours during pregnancy: A systematic review. Int J Behav Nutr Phys Act

- 523 2017;14:32.
- 524 36. Oviedo-Caro MA, Bueno-Antequera J, Munguía-Izquierdo D. Measuring  
525 Sedentary Behavior During Pregnancy: Comparison Between Self-reported and  
526 Objective Measures. *Matern Child Health J* 2018;22:968–977.
- 527 37. Ruifrok AE, Althuisen E, Oostdam N, Van Mechelen W, Mol BW, De Groot  
528 CJM, Van Poppel MNM. The relationship of objectively measured physical  
529 activity and sedentary behaviour with gestational weight gain and birth weight. *J*  
530 *Pregnancy* 2014;2014:567379.
- 531 38. Melzer K, Schutz Y, Boulvain M, Kayser B. Pregnancy-related changes in  
532 activity energy expenditure and resting metabolic rate in Switzerland. *Eur J Clin*  
533 *Nutr* 2009;63:1185–1191.
- 534 39. Oviedo-Caro M, Bueno-Antequera J, Munguía-Izquierdo D. Transcultural  
535 adaptation and psychometric properties of Spanish version of Pregnancy Physical  
536 Activity Questionnaire: the PregnActive project. *Gac Sanit* 2018;
- 537 40. Price BB, Amini SB, Kappeler K. Exercise in pregnancy: Effect on fitness and  
538 obstetric outcomes - A randomized trial. *Med Sci Sports Exerc* 2012;44:2263 –  
539 2269.
- 540 41. Phelan S. Pregnancy: a ‘teachable moment’ for weight control and obesity  
541 prevention. *Am J Obstet Gynecol* 2010;202:e1–e8.
- 542 42. Baena-García L, Ocón-Hernández O, Acosta-Manzano P, Coll-Risco I, Borges-  
543 Cosic M, Romero-Gallardo L, de la Flor-Alemanly M, Aparicio VA. Association  
544 of sedentary time and physical activity during pregnancy with maternal and  
545 neonatal birth outcomes. The GESTAFIT Project. *Scand J Med Sci Sport*  
546 2019;29:407–414.

**Table 1.** Associations between activity composition<sup>a</sup> and adiposity and CRF: Results from Wald chi square type II ANOVA tests of linear models.

	Sum Sq	Df	F value	Pr(>F)
Sum of skinfold thickness, mm.†	2.6	3	13.9	0.000
FMI, kg/m <sup>2</sup> .†	4.9	3	17.1	0.000
BMI, kg/m <sup>2</sup> .†	0.9	3	14.8	0.000
CRF (VO <sub>2max</sub> ), mL/kg·min.	906.6	3	20.3	0.000
CRF (VO <sub>2max</sub> ), mL/kg·min. Additionally adjusted for sum of skinfold thickness.	175.5	3	5.7	0.001
CRF (VO <sub>2max</sub> ), mL/kg·min. Additionally adjusted for FMI.	49.2	3	3.2	0.025
CRF (VO <sub>2max</sub> ), mL/kg·min. Additionally adjusted for BMI.	111.5	3	6.2	0.001

Abbreviations: FMI: fat mass index; BMI: body mass index; CRF: cardiorespiratory fitness; VO2max: maximal oxygen consumption. <sup>a</sup> Activity composition expressed as isometric log-ratios. †Log transformed variable. All models are adjusted for age, educational level and employment status.

**Table 2.** Estimated percentage of the differences from the mean composition in predicted adiposity associated with 15-minutes and 30-minutes reallocations.

<i>15-minutes reallocation</i>					<i>30-minutes reallocation</i>				
<b>Sum of skinfold thickness, mm</b>									
	Sleep	ST	LPA	MVPA		Sleep	ST	LPA	MVPA
Sleep		-0.3	<b>1.5</b>	<b>2.4</b>	Sleep		-0.6	<b>3.1</b>	<b>5.5</b>
ST	0.2		<b>1.8</b>	<b>2.7</b>	ST	0.5		<b>3.6</b>	<b>6.0</b>
LPA	<b>-1.4</b>	<b>-1.7</b>		1.0	LPA	<b>-2.8</b>	<b>-3.3</b>		2.5
MVPA	<b>-2.0</b>	<b>-2.3</b>	-0.5		MVPA	<b>-3.8</b>	<b>-4.2</b>	-0.8	
<b>FMI, kg/m<sup>2</sup></b>									
	Sleep	ST	LPA	MVPA		Sleep	ST	LPA	MVPA
Sleep		-0.8	<b>1.7</b>	<b>3.1</b>	Sleep		-1.6	<b>3.5</b>	<b>7.1</b>
ST	0.8		<b>2.5</b>	<b>3.9</b>	ST	1.5		<b>5.0</b>	<b>8.7</b>
LPA	<b>-1.6</b>	<b>-2.4</b>		1.5	LPA	<b>-3.1</b>	<b>-4.6</b>		3.8
MVPA	<b>-2.5</b>	<b>-3.3</b>	-0.9		MVPA	<b>-4.6</b>	<b>-6.1</b>	-1.3	
<b>BMI, kg/m<sup>2</sup></b>									
	Sleep	ST	LPA	MVPA		Sleep	ST	LPA	MVPA
Sleep		-0.5	0.5	1.2	Sleep		-1.1	1.0	<b>2.7</b>
ST	0.5		<b>1.0</b>	<b>1.7</b>	ST	1.1		<b>2.1</b>	<b>3.8</b>
LPA	-0.5	<b>-1.0</b>		0.7	LPA	-0.9	<b>-2.0</b>		1.8
MVPA	-0.9	<b>-1.5</b>	-0.5		MVPA	-1.7	<b>-2.8</b>	-0.7	

Reallocation of 15 and 30 minutes are from the activity in the columns to the activity in the rows. Values are expressed in % of the differences from the mean composition in predicted adiposity. Bold type indicates significant change in outcome variable. Abbreviations: FMI: fat mass index; BMI: body mass index; ST: sedentary time; LPA: light physical activity; MVPA: moderate-to-vigorous physical activity. Analysis adjusted for age, education level, and employment status.



**Table 3.** Estimated percentage of the differences from the mean composition in predicted CRF associated with 15-minutes and 30-minutes reallocations.

<i>15-minutes reallocation</i>					<i>30-minutes reallocation</i>				
<b>CRF, mL/kg·min *</b>									
	Sleep	ST	LPA	MVPA		Sleep	ST	LPA	MVPA
Sleep		0.1	<b>-0.7</b>	<b>-1.8</b>	Sleep		0.1	<b>-1.5</b>	<b>-4.0</b>
ST	0.0		<b>-0.8</b>	<b>-1.8</b>	ST	-0.1		<b>-1.6</b>	<b>-4.1</b>
LPA	<b>0.7</b>	<b>0.8</b>		<b>-1.1</b>	LPA	<b>1.4</b>	<b>1.5</b>		<b>-2.6</b>
MVPA	<b>1.5</b>	<b>1.6</b>	0.8		MVPA	<b>2.9</b>	<b>3.0</b>	1.4	
<b>CRF, mL/kg·min (Additionally adjusted for the sum of skinfold thickness)</b>									
	Sleep	ST	LPA	MVPA		Sleep	ST	LPA	MVPA
Sleep		0.1	-0.3	<b>-1.0</b>	Sleep		0.1	-0.6	<b>-2.2</b>
ST	0.0		-0.3	<b>-1.0</b>	ST	-0.1		-0.7	<b>-2.3</b>
LPA	0.3	0.3		-0.7	LPA	0.6	0.7		-1.6
MVPA	<b>0.8</b>	<b>0.9</b>	0.5		MVPA	<b>1.6</b>	<b>1.7</b>	0.9	
<b>CRF, mL/kg·min (Additionally adjusted for FMI)</b>									
	Sleep	ST	LPA	MVPA		Sleep	ST	LPA	MVPA
Sleep		-0.2	-0.2	<b>-0.8</b>	Sleep		-0.4	-0.4	<b>-1.7</b>
ST	0.2		0.0	<b>-0.6</b>	ST	0.4		-0.1	<b>-1.4</b>
LPA	0.2	0.0		<b>-0.6</b>	LPA	0.5	0.1		<b>-1.3</b>
MVPA	<b>0.7</b>	<b>0.5</b>	0.5		MVPA	<b>1.3</b>	<b>0.9</b>	0.8	
<b>CRF, mL/kg·min (Additionally adjusted for BMI)</b>									
	Sleep	ST	LPA	MVPA		Sleep	ST	LPA	MVPA
Sleep		-0.3	<b>-0.4</b>	<b>-1.0</b>	Sleep		-0.6	<b>-0.8</b>	<b>-2.3</b>
ST	0.3		-0.1	<b>-0.8</b>	ST	0.6		-0.2	<b>-1.7</b>
LPA	<b>0.4</b>	0.1		<b>-0.6</b>	LPA	<b>0.8</b>	0.2		<b>-1.5</b>
MVPA	<b>0.9</b>	<b>0.6</b>	0.5		MVPA	<b>1.7</b>	<b>1.1</b>	0.9	

Reallocation of 15 and 30 minutes are from the activity in the columns to the activity in the rows. Values are expressed in % of the differences from the mean composition in predicted CRF. Bold type indicates significant change in outcome variable.

Abbreviations: CRF: cardiorespiratory fitness; FMI: fat mass index; BMI: body mass index; ST: sedentary time; LPA: light physical activity; MVPA: moderate-to-vigorous physical activity. \*Analysis adjusted for age, education level, and employment status.